Very Long Baseline Oscillations

The BNL VLBL Concept

Brett Viren

Physics Department

BROOKHAVEN NATIONAL LABORATORY

7th International Workshop on Neutrino Factories and Superbeams

Outline

1. The Concept
2. Oscillation Probability Illustrations
3. Example: BNL to Homestake
4. Making Very Long Baseline Neutrinos at BNL

1 The Concept
- Goals and Overview
- Motivations for Basic Parameters
- Natural Benefits to Going High Energy

2 Oscillation Probability Illustrations

3 Example: BNL to Homestake

4 Making Very Long Baseline Neutrinos at BNL

BROOKHAVEN NATIONAL LABORATORY

Basics of the VLBL Concept

The VLBL concept consists of three simple ideas:

1. Use a very **long** baseline,
2. a **wide** band ν-beam,
3. at **high** ν-energies.

This allows an ambitious but affordable experiment which is qualitatively different than previous or planned LBL experiments.

- It is sensitive to multiple physical effects.
- Allows one to break parameter degeneracies.
- Rough measurement of δ_{CP} and θ_{13} with only ν-running.

If we get lucky with the true parameter values, these statements can be made more strongly.

BROOKHAVEN NATIONAL LABORATORY

Why Long, Wide and High?



1. Wide → see multiple oscillations
2. High → avoid Fermi motion
3. High ∴ Long

Blue bars show flux coverage (full width at 10% peak)

BROOKHAVEN NATIONAL LABORATORY

Natural Background Reduction





Putting some $\nu_\mu \rightarrow \nu_e$ signal at high energy, gain strong suppression of NC bkg via Q^2 kinematic cutoff

- Nuance MC
- Q^2 (top) and E_{π^0} (bottom) for single-π NC events
- Each color band: mono-energetic neutrinos, 1-10 GeV in 1 GeV steps.
- At $E_\nu > 2$ GeV get $> 50\times$ natural background reduction

1 The Concept

2 Oscillation Probability Illustrations
- Disappearance
- Appearance

3 Example: BNL to Homestake

4 Making Very Long Baseline Neutrinos at BNL

Oscillation Probability Illustrations

Unless noted, the values used are:

Δm^2_{21} = 8.0e-5 eV^2. KamLAND/SK/SNO.

Δm^2_{32} = 2.5e-3 eV^2. Super-Kamiokande.

$\sin^2(2\theta_{23})$ = 1.0. SK atmospheric maximal mixing.

$\sin^2(2\theta_{12})$ = 0.86. SK/SNO solar mixing.

$\sin^2(2\theta_{13})$ = 0.04. CHOOZ limited: use "not too big, not too small".

δ_{CP} = 0. Totally unknown.

Baseline = 2540 km (other, short baselines shown for comparison)

Matter effects using PREM Earth density and electron fraction profile.

Probability plots:

- Energy range will reflect full-width-10%-max coverage.
- Flash short baseline examples to compare.

Multiple Disappearance Oscillations Well Covered!



- Multiple wiggles!
- No problem resolving "the dip", will see more than one!
- Nodes well placed across flux coverage, robust against Δm^2 change.

Disappearance at Short Baselines

At most one dip to see. Can slip away if Δm^2 ends up low.





Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

- Multiple Appearance Peaks

Multiple Appearance Peaks Covered



- Multiple peaks covered
- Robust against Δm^2 change

Appearance at Short Baselines





BROOKHAVEN NATIONAL LABORATORY

Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

- Multiple Appearance Peaks
- Matter Effects

Matter Effects at Very Long Baselines



Sensitivity to $sign(\Delta m^2)$

- Factor of 2 effect → "easy"
- Effect mostly in first peak $E_\nu > 2$ GeV

Matter Effects at Short Baselines





Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

- Multiple Appearance Peaks
- Matter Effects
- CP Violation

Nonzero CP Angle at Very Long Baselines



- Shift in magnitude and location
- Stronger effect for oscil peak $n > 1$ (W.Marciano)
- Longer baseline pulls $n > 1$ away from Fermi-motion region.

Nonzero CP Angle at Short Baselines





Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

- Multiple Appearance Peaks
- Matter Effects
- CP Violation
- ν_e Appearance with $\theta_{13} = 0$!

Guaranteed ν_e Appearance



What if $\theta_{13} = 0$?
Still observe ν_e via Δm^2_{21} !

- $\theta_{13} = 0$ in black
- $\theta_{13} \neq 0$ in blue
- Upswing at low energies is due to Δm^2_{21}.
- Small shift if non-maximal mixing

This over constrains the solar parameters - potential to see new physics!

$\theta_{13} = 0$ at Short Baselines





Rich Appearance Effects

The Very Long Baseline approach makes available a rich set of appearance effects.

- Multiple Appearance Peaks
- Matter Effects
- CP Violation
- ν_e Appearance with $\theta_{13} = 0$!

Since these effects occur differently across the spectrum in a Very Long Baseline experiment, they can be disentangled with a single experiment.

Sensitivity to Different Parameters in Different Energy Regions

	$E_\nu < 1$ GeV	$1 < E_\nu 2 <$GeV	$E_\nu > 2$ GeV
$\sin^2 2\theta_{13}$	$\surd$	$\surd$	$\surd$
$sign(\Delta m^2_{32})$	-	-	$\surd\surd\surd$
δ_{CP}	$\surd$	$\surd\surd$	$\surd$
solar	$\surd\surd\surd$	$\surd$	-

- It's a complex picture with many effects!
- But, effects have different strength at different energies.
- Measuring across the wide energy band makes it possible to sort them out.

1 The Concept

2 Oscillation Probability Illustrations

3 Example: BNL to Homestake
- Event Rate
- Disappearance
- Appearance

4 Making Very Long Baseline Neutrinos at BNL

BROOKHAVEN NATIONAL LABORATORY

Example Baseline: 2540 km



Homestake & Henderson equivalent. Assume UNO class far detector.

Neutrino Flux

The following is work by Milind Diwan.



ν running:

- 1 MW, 28 GeV proton beam
- 5×10^7 seconds
- 1.12×10^{22} PoT
- 60 cm carbon target
- 4 m $\phi\times$ 200 m long decay tunnel

$\bar{\nu}$ running same but 2 MW

Expected Number of Events

- 1 MW ν running
- Water-Cherenkov detector
- 500 kTon water fiducial
- 2540 km baseline
- 5×10^7 seconds exposure

Reaction	Number
CC $\nu_\mu + N \rightarrow \mu^- + X$	51800
NC $\nu_\mu + N \rightarrow \nu_\mu + X$	16908
CC $\nu_e + N \rightarrow e^- + X$	380
QE $\nu_\mu + n \rightarrow \mu^- + p$	11767
QE $\nu_e + n \rightarrow e^- + p$	84
CC $\nu_\mu + N \rightarrow \mu^- + \pi^+ + N$	14574
NC $\nu_\mu + N \rightarrow \nu_\mu + N + \pi^0$	3178
NC $\nu_\mu + O^{16} \rightarrow \nu_\mu + O^{16} + \pi^0$	574
CC $\nu_\tau + N \rightarrow \tau^- + X$ (if all $\nu_\mu \rightarrow \nu_\tau$)	319

BROOKHAVEN NATIONAL LABORATORY

Disappearance at 2540 km



- 10% energy resolution + Fermi motion
- 2 nodes clearly visible
- **Black** and **red** include background
- Only single-π production bkg considered here
- $\sigma(\Delta m^2_{32})$ & $\sigma(\sin^2(2\theta_{23})) \sim 1\%$
- Resolution dominated by systematics

Expected Appearance Results

1. Weighted SK Atm-ν MC/reconstruction study (C. Yanagisawa)
2. Baseline requirements study (M. Diwan)
3. In progress: full study with UNO detector MC and realistic reconstruction code.

C. Yanagisawa - weighted SK Atm-ν MC study

Complete water Cherenkov detector simulations progress

ν_e CC for signal ; all $\nu_{\mu,\tau,e}$ NC , ν_e beam for background

- Δm^2_{21}=7.3 x 10^{-5} eV^2, Δm^2_{31}=2.5 x 10^{-3}eV^2
- $\sin^2 2\theta_{ij}$(12,23,13)=0.86/1.0/0.04, δ_{CP}=+45,+135,-45,-135°





Select single ring events and select electrons
Signal/backg = 700/2005

Perform analysis of single electron pattern, likelihood cut retaining ~50% of signal.
Signal/back = 321/169

C. Yanagisawa (Stony Brook), 3rd BNL/UCLA workshop
http://www.physics.ucla.edu/hep/proton/proton2005.htm

Improved π^0 finder + likelihood cut.

Appearance - Baseline Requirements Study

M. Diwan, close but stricter requirements still:





Sensitivity of δ_{CP} and θ_13

1-σ and 90% contours, all other parameters fixed.





BROOKHAVEN NATIONAL LABORATORY

Sensitivity of δ_{CP} and θ_13 - Combined $\nu/\bar{\nu}$

1-σ contours, all other parameters fixed.

Regular hierarchy νυ and Antiνυ running



Reversed hierarchy νυ and Antiνυ running



Patrick Huber, UWisc:



- Both ν- and $\bar{\nu}$-running.
- Includes correlations and errors on all parameters, 10% background uncertainty.
- Will improve as other oscillation parameter measurements improve.
- All δ_{CP} range covered.

1 The Concept

2 Oscillation Probability Illustrations

3 Example: BNL to Homestake

4 Making Very Long Baseline Neutrinos at BNL
- AGS Upgrade
- BNL Site Development
- Target and Horn
- Price Tag

BROOKHAVEN NATIONAL LABORATORY

Upgrade to AGS



- Use 1.2 GeV SC Linac instead of booster
  - $7 \times 10^{13} \rightarrow 9 \times 10^{13}$ ppp
  - Fill time 0.6 sec $\rightarrow$ 1.0 msec
- Increase rep. rate 0.5 $\rightarrow$ 2.5 Hz
- New mag. PS and RF cavities
- Further improvements on design being worked on.

BNL Site Development





- Point to most points west
- Keep hadrons above the water table
- Room for a (very) near detector
- Hill cheaper than tunnel

BROOKHAVEN NATIONAL LABORATORY



- Conventional pulsed hadron focussing with 2 horns
- Likely carbon-carbon target, embedded in 1st horn
- R&D underway with material experiments
- Collaboration with FNAL, JPARC and others

Estimated Costs

pgflastimage

- Put through internal BNL review
- Bottom-up estimation with WBS
- Based on RHIC & SNS ring, LHC magnets.
- Target & Horn from BNL, K2K and NuMI.
- $273.4M FY04
- 6 Years to neutrinos
  - 3 years R&D
  - Construction after 1 year
  - 4.5 years to completion
  - 0.5 year commissioning

BROOKHAVEN NATIONAL LABORATORY

Summary

- The BNL Very Long Baseline concept provides a qualitatively different experiment than past or proposed
- Precision (1%) measurement of atm-ν params, systematics limited.
- Degeneracy-broken measurement of appearance parameters
- If lucky, ν-running only needed for appearance measurements, with $\bar{\nu}$-running gain precision w/out needing luck.
- Appearance results limited by how well background can be controlled
- Work on full detector simulation and reconstruction to be finished before final word.
- Affordable, practical beam design. No "magic" needed.
- Ongoing collaborations (C. Yanagisawa (UNO), P. Huber (GLoBES)), others most welcome!

BROOKHAVEN NATIONAL LABORATORY

References

Papers:

- BNL Neutrino Working Group web page: http://nwg.phy.bnl.gov/.
- "Extra Long Baseline Neutrino Oscillations and CP Violation", W. Marciano, BNL-HET-01/31, hep-ph/0108181.
- Whitepaper, BNL-69395 hep-ex/0211001.
- VLBL PRD paper, Phys.Rev. D68 (2003) 012002, hep-ph/0303081.
- AGS Superbeam CDR, BNL-73210-2004-IR (avail from NWG web).
- "The Case for a Super Neutrino Beam", M. Diwan (1290 km and 2540 km compared) hep-ex/0407047.

Publicly available software:

- Oscillation Probability Calculator, B. Viren, http://minos.phy.bnl.gov/ bviren/elbo/libnuosc++/
- GLoBES, P. Huber, M. Lindner and W. Winter, hep-ph/0407333
- Nuance, D. Casper, hep-ph/0208030

BROOKHAVEN NATIONAL LABORATORY

Backup Slides

CP Violation Sensitivity Independent of Baseline



- Marciano, hep-ph/0108181
- $A \sim L$
- Flux $\sim 1/L^2$
- Statistical F.O.M.
  - $= (\Delta A/A)^{-2}$
    $= A^2 N/(1-A)$
  - $N = N_{\nu e} + N_{\bar{\nu} e}$
  - Linear in flux
  - Quadratic in A

Independent of BL, at least until the flux totally runs out.

Resolution on θ_{23} and Δm^2_{32}



- Energy scale systematic not included (SK has 2.5%)
- O.w. $\sigma(\Delta m^2_{32}) \sim 1\%$
- Not strongly sensitive on normalization uncertainty due to multi-nodal spectrum shape.

BROOKHAVEN NATIONAL LABORATORY